### USE OF EXTRACTS OF CINNAMON (*Cinnamomum zeylanicum Blume*) AS DYEING AGENT ON COTTON FABRIC

BY

#### W. D. I. U. JAYARATHNE



# FACULTY OF TECHNOLOGY EASTERN UNIVERSITY

#### SRI LANKA

2023

# **TABLE OF CONTENT**

ABSTRACTi
ACKNOWLEDGEMENTS ii
LIST OF TABLES vii
LIST OF FIGURES viii
ABBREVIATIONSix
CHAPTER 011
1.0. INTRODUCTION
1.1. Background1
1.2. Objectives
CHAPTER 02
2.0. LITERATURE REVIEW
2.1. Textile dyeing
2.2. Nature of fabric
2.2.1. Cellulose
2.2.2. Cotton
2.2.3. Chemical composition of Cotton
2.3. Dyes
2.3.1. History of Dyes
2.4. Different types of dye

2.4.1. Synthetic dye
2.4.2. Natural dye
2.4.2.1. Based on chemical structure
2.4.2.2. Based on the sources
2.4.2.3. Based on the application method
2.5. Dye extraction
2.5.1. Aqueous extraction
2.5.2. Acid and alkali extraction
2.5.3. Solvent extraction
2.6. Principle of natural dyeing
2.6.1. Pre-treatment
2.6.2. Mordants
2.6.2.1. Metallic mordants
2.6.2.1.1. Brightening mordants
2.6.2.1.2. Dulling mordants
2.6.2.2. Oils type mordants
2.6.2.3. Tannins
2.6.3. Mordanting
2.6.3.1. Pre-mordanting
2.6.3.2. Meta-mordanting/simultaneous mordanting17

2.6.3.3. Post-mordanting
2.7. Cinnamon
2.7.1. Cinnamon in Sri Lanka
2.7.2. Cinnamomum Zeylanicum Blume (Ceylon cinnamon)
2.7.3. Chemical constituent of Ceylon cinnamon19
2.7.3.1. Cinnamaldehyde19
2.7.4. Anti-bacterial properties of cinnamon
CHAPTER 03
3.0. MATERIALS AND METHODOLOGY
3.1. Materials
3.1.1. Samples collection
3.1.2. Chemicals
3.1.3. Imaging
3.2. Experiment
3.2.1. Pre-treatment of cotton source
3.2.2. Oxidation of cotton using H <sub>2</sub> O <sub>2</sub>
3.2.3. Determination of carboxyl acid content of cotton25
3.2.4. Extraction of cinnamon dye
3.2.5. Mordanting step
3.2.6. Dye immerse step

3.3. Data collection
3.3.1. Anti-bacterial efficacy of the dye-fixed-fabric
3.3.1.1. Preparation of nutrient agar
CHAPTER 04
4.0. RESULTS AND DISCUSSION
4.1. Oxidation of cotton using H <sub>2</sub> O <sub>2</sub>
4.2. Determination of carboxyl acid content of cotton
4.3. Results of dyeing and color intensity
4.4. Anti-bacterial efficacy of the dye and dye-fixed-fabric
CHAPTER 05
REFERENCES

## **LIST OF TABLES**

Table 3.1: Chemicals amounts of pre-treatment solution	23
Table 3.2: Chemicals amounts of oxidation solution	25
Table 3.3: Composition of Nutrition agar	29
Table 4.1: Weight of cotton fabric pre and post oxidation	31
Table 4.2: Titration results of acetic acid neutralization by NaOH	32

### **LIST OF FIGURES**

Figure 2.1: Cellulose structure
Figure 2.2: Natural colorant classification
Figure 2.3: Cellulose, mordant metal ion and natural dye complex
Figure 2.4: Cinnamaldehyde
Figure 3.1: Pre-treatment of cotton fabric
Figure 3.2: Extraction of cinnamon dye
Figure 3.3: Mordanting
Figure 4.1: Color produced by cinnamon dye
Figure 4.2: Testing anti-bacterial efficacy of control sample
Figure 4.3: Testing anti-bacterial efficacy of dye and dye fixed fabric sample

#### ABSTRACT

Natural dyeing is considered as a suitable approach in textile dyeing, which can minimize the environmental impacts of synthetic dyes. The present study has shown that it is possible to use the bark of *Cinnamomum zyelanicum Blume* (cinnamon) to extract natural dyes using aqueous extraction method for coloring 100% cotton fabric. Cinnamaldehyde was the agent of interest for the current project as a dyeing agent. In order to fix cinnamaldehyde on cotton material, metal salts of Cu, Ni, Fe, Al, and Co were used during experiments. According to the results, a darker shade was seen with FeSO4-mordanted cotton fabric. A CuSO<sub>4</sub>-mordanted generated a medium shade. In the cotton samples mordanted with NiCl<sub>2</sub>, CoCl<sub>2</sub>, and (NH<sub>4</sub>)Al(SO<sub>4</sub>)<sub>2</sub>, relatively light colors were seen.

Moreover, the antibacterial activity of dye and dye-fixed fabric against textile bacteria was investigated. The inhibitory effect was observed using the disk diffusion method. Cinnamon dye and cinnamon-dyed cotton fabric were not active against the tested bacteria culture.

î

Key words: Cinnamomum zeylanicum, Cinnamaldehyde; Antibacterial activity