

In vitro studies on plant development of *Camellia sinensis* (L.) O.
Kuntze (tea) with emphasis on somatic embryogenesis



Thayamini Harold Seran



ExPhD16



Thesis
Library - EUSL

Thesis submitted for the degree of **Doctor of Philosophy**

62846

Department of Plant Sciences

Faculty of Science

University of Colombo

Sri Lanka

2006

PROCESSED
EUSL Library, EUSL

Abstract

Histological and morphological studies on somatic embryogenesis were carried out for efficient *in vitro* plant development of *Camellia sinensis* (tea). To determine the suitable stage of cotyledons for direct somatic embryogenesis, the sterilized cotyledons (6 -12 mm diameter) at various developmental stages were inoculated on MS medium containing 2.0 mg L⁻¹ BAP and 0.2 mg L⁻¹ NAA. Results indicated that immature cotyledon (10 mm diameter, stage 3) was the most responsive stage for the production of typical (normal) somatic embryos. Cytological examinations revealed that typical somatic embryos at different stages exhibit gradual tissue differentiations. Typical mature somatic and zygotic embryos morphologically and histologically have similar bipolar structures.

Immature cotyledon segments (stage 3) were placed on different media containing NAA (0.0 – 3.0 mg L⁻¹) in combination with BAP (2.0 mg L⁻¹) to select suitable medium for efficient production of typical somatic embryos. The optimum concentrations of NAA were in the range of 0.2 – 1.0 mg L⁻¹ for direct somatic embryogenesis (14.6% - 16.7%). Cotyledon-type and seed-like somatic embryos were developed in 0.2 and 1.0 mg L⁻¹ NAA respectively. Further, the explants of high yielding cultivars DT1 (Estate selection in Sri Lanka) and TRI 2025 (Assam type) performed well and exhibited embryogenic potential (28.1% and 21.9% respectively).

Several types of explants obtained from *in vitro* germinated zygotic embryonic axes were cultured to select the suitable explants for the production of typical somatic embryos. Results indicated that both hypocotyl and small succulent leaf explants were

the best for the production of firm, typical somatic embryos. Histological studies showed that most somatic embryos were directly originated from the cortex tissue of the hypocotyl and upper epidermal layer of the succulent leaf explants.

Further, attempts were made to establish *in vitro* callus culture from field grown leaves. Results indicated that high frequencies (50% - 58 %) of aseptic cultures were achieved in 60% and 75% solutions of Clorox™ for 30 min soaking period among varying concentrations of disinfectant (15% - 75%) and various exposure times (15 - 60 min). The rates of initiation (72.2%) and formation (56.9%) of calli from *in vitro* aseptic explants were relatively high in MS medium containing BAP (2.0 mg L⁻¹) and NAA (3.0 mg L⁻¹) among tested combinations of BAP and NAA concentrations.

Morphological and histological studies were carried out on the development of embryogenic callus from leaves of *in vitro* shoots. It was observed that collenchyma cells were developed from vegetative cells at the cut end and packed within the swelling tissues. After bursting, parenchyma cells were gradually formed as a result of cell divisions. These parenchyma cells differentiated as embryogenic cells. The average number of single embryogenic cells was significantly higher in the liquid than in solid medium. Further investigations were done to produce somatic embryos from primary embryogenic calli. Results showed that formation of somatic embryos (11.7%) was high in MS medium containing BAP (1.0 mg L⁻¹) and NAA (0.1 mg L⁻¹) in combination with GA₃ (0.1 mg L⁻¹).

Zygotic embryonic axes were coated with 3% and 4% sodium alginate matrices. The results revealed that 3% sodium alginate with 100 mM calcium chloride dihydrate provided the most suitable matrix concentrations for encapsulation. Further, naked and coated axes cultured on MS medium without (MS1) or with (MS2) 3.0 mg L⁻¹ BAP and 0.5 mg L⁻¹ IBA gave highest rates of survival (100%) and germination (93-100%). But efficient plant recovery from non-encapsulated (73.3%) and encapsulated (42.2%) axes was achieved on MS2 medium. A high rate (48%) of *in vitro* plantlets was survived at 8th week of transplanting in soil than in a mixture of soil: sand (1:1 v/v).

Feasibility of cold storage of encapsulated embryonic axes was studied to obtain efficient *in vitro* conversion of plants and their survival under *ex vitro* conditions. Synthetic seeds and natural seeds were stored at 4 °C for 0, 4, 8, 16, and 20 weeks. Efficient *in vitro* germination and plant recovery were obtained from synthetic seeds stored for four weeks than control. Vigorous plantlets were collected from *in vitro* cultures and successfully established in soil: coconut coir dust (1:1 v/v).

To perform synchronous germination of plant material into vigorous plants under *ex vitro* conditions, sterilized zygotic embryos and zygotic embryonic axes were cultured on germination medium whereas seeds were sown in sand bed as a control. Consistent germination was observed on cultured embryonic axes *in vitro* for uniform growth where rate of germination was significantly highest (99.0%) at 4th week. Further, healthy plantlets developed from embryonic axes *in vitro* had erect shoots with short internodes as well as taproots with abundance adventitious roots under nursery conditions for better adaptation in field as compared with seedling raised by conventional sexual propagation.

Contents

	page
Acknowledgements	i
Abstract	iii
Contents	vi
Abbreviations	ix
List of Figures	x
List of Tables	xi
Chapter 1: General Introduction	1
1.1 Tea and its importance	1
1.2 Conventional breeding and propagation	3
1.3 <i>In vitro</i> propagation techniques	4
1.4 <i>In vitro</i> somatic embryogenesis	6
1.5 <i>In vitro</i> storage via encapsulation	9
1.6 <i>In vitro</i> plant regeneration	10
1.7 Performance of <i>in vitro</i> plantlets under <i>ex vitro</i> conditions	11
1.8 Objectives of the study	12
Chapter 2: Morphological and histological studies on somatic embryogenesis from cotyledon explants of <i>Camellia sinensis</i> (L.)	13
2.1 Introduction	13
2.2 Materials and Methods	15
2.2.1 <i>In vitro</i> somatic embryogenesis	15
2.2.1.1 Effect of development stages of intact cotyledons	15
2.2.1.2 Sample preparations for histological studies	16
2.2.1.3 Effect of NAA on cotyledon segments	17
2.2.1.4. Effect of cultivars on cotyledonary tissues	17
2.2.2 Culture conditions for <i>in vitro</i> germination	18
2.2.3 Statistical analysis	19
2.3 Results	19
2.3.1 <i>In vitro</i> somatic embryogenesis	19
2.3.1.1 Effect of development stages of intact cotyledons	19
2.3.1.2 Histology of somatic and zygotic embryos	21
2.3.1.3 Effect of NAA on cotyledon segments	25
2.3.1.4 Effect of cultivars on cotyledonary tissue	25
2.3.2 Germination of <i>in vitro</i> somatic embryos to plantlets	27
2.4 Discussion	30
Chapter 3: Direct somatic embryogenesis from explants obtained from <i>in vitro</i> germinated embryonic axes of <i>Camellia sinensis</i> (L.)	38
3.1 Introduction	38
3.2 Materials and methods	40
3.2.1 Source of explants	40
3.2.2 <i>In vitro</i> somatic embryogenesis	40

3.2.3	Cytological examination of explants in culture	42
3.2.4	Culture conditions for <i>in vitro</i> germination of somatic embryos	42
3.2.5	Statistical analysis	43
3.3	Results	43
3.3.1	<i>In vitro</i> germination of zygotic embryonic axes	43
3.3.2	<i>In vitro</i> somatic embryogenesis	43
3.3.3	Histology changes in the explants during somatic embryogenesis	49
3.3.4	Germination of <i>in vitro</i> somatic embryos to plantlets	52
3.4.	Discussion	52

Chapter 4: Indirect somatic embryogenesis from leaf explants of *Camellia sinensis* (L.) 57

4.1	Introduction	57
4.2	Materials and methods	59
4.2.1	Establishment of <i>in vitro</i> callus from leaf explants	59
4.2.1.1	Culture conditions for initiation of callus	59
	<i>Surface sterilization of leaves</i>	59
	<i>Initiation of callus from field grown leaves</i>	60
	<i>Initiation of callus from in vitro leaves</i>	61
4.2.1.2	Source of leaf explants for efficient callus formation	62
4.2.2	Production of embryogenic callus from <i>in vitro</i> leaves	63
4.2.2.1	Culture conditions for initiation and maintenance of embryogenic callus	63
4.2.2.2	Histological studies on the development of embryogenic callus	63
4.2.2.3	Effect of solid and liquid media on intensity of embryogenic cells	64
4.2.3	<i>In vitro</i> somatic embryogenesis from embryogenic callus	65
4.2.3.1.	Effect of the strength of MS basal medium and ABA	65
4.2.3.2.	Effect of culture duration	65
4.2.3.3.	Effect of some selected combinations of growth regulators	68
4.2.4	Statistical analysis	68
4.3	Results and Discussion	69
4.3.1	Establishment of <i>in vitro</i> callus culture from leaves	69
4.3.1.1	Initiation of callus from leaf explants	69
	<i>Surface sterilization of leaves</i>	69
	<i>Callus initiation from field grow leaves</i>	75
	<i>Callus initiation from in vitro leaves</i>	75
4.3.1.2	Efficient callus formation	79
4.3.2	Production of embryogenic callus from <i>in vitro</i> leaves	81
4.3.2.1	Initiation and maintenance of embryogenic callus	81
4.3.2.2	Histology of embryogenic callus	84
4.3.2.3	Effect of solid and liquid media on intensity of embryogenic cells	88
4.3.3	<i>In vitro</i> somatic embryogenesis from embryogenic-callus	90
4.3.3.1	Effect of the strength of MS basal medium and ABA	90
4.3.3.2	Effect of culture duration	93
4.3.3.3	Effect of some selected combinations of growth regulators	94

Chapter 5: Cold storage of encapsulated zygotic embryonic axes of <i>Camellia sinensis</i> (L.)	98
5.1 Introduction	98
5.2 Materials and methods	100
5.2.1 <i>In vitro</i> germination of synthetic seeds	100
5.2.1.1 Plant material	100
5.2.1.2 Effect of encapsulating matrix on the production of synthetic seeds	100
5.2.1.3 <i>In vitro</i> germination of encapsulated embryonic axes	101
5.2.1.4 Establishment of <i>In vitro</i> plantlets under <i>ex vitro</i> conditions	102
5.2.1.5 Statistical analysis	103
5.2.2 <i>In vitro</i> storage of synthetic seeds	103
5.2.2.1. Source of material	103
5.2.2.2 Encapsulation of zygotic embryonic axes	103
5.2.2.3 Storage of synthetic and natural seeds	104
5.2.2.4 Culture conditions for <i>in vitro</i> germination of stored materials	104
5.2.2.5 Acclimatization and nursery establishment of <i>in vitro</i> plantlets	105
5.2.2.6 Statistical analysis	106
5.3 Results	106
5.3.1 <i>In vitro</i> germination of synthetic seeds	106
5.3.1.1 Encapsulation of zygotic embryonic axes	106
5.3.1.2 <i>In vitro</i> germination of encapsulated embryonic axes	107
5.3.1.3 Performance of <i>in vitro</i> plantlets under <i>ex vitro</i> conditions	111
5.3.2 <i>In vitro</i> storage of synthetic and natural seeds	111
5.3.2.1. Viability and <i>in vitro</i> germination of stored seeds	111
5.3.2.2 Plant development of stored material	113
5.3.2.3 Establishment of <i>in vitro</i> plantlets under <i>ex vitro</i> conditions	113
5.4 Discussion	117
Chapter 6: Germination capacity of <i>in vitro</i> zygotic embryonic axes of in comparison with conventional seeds of <i>Camellia sinensis</i> (L.) and their subsequent plant development	126
6.1 Introduction	126
6.2 Materials and Methods	128
6.2.1 Plant material	128
6.2.2 Culture conditions for seedling and <i>in vitro</i> plantlet development	128
6.2.3 Establishment of seedling and <i>in vitro</i> plantlet under <i>ex vitro</i> conditions	128
6.2.4 Statistical analysis	129
6.3 Results	129
6.3.1 Germination of plant material under <i>in vitro</i> and <i>ex vitro</i> conditions	129
6.3.2 Plant growth and morphology under <i>ex vitro</i> conditions	131
6.4 Discussion	137
Chapter 7: General Discussion	142
Chapter 8: References	156
Appendices	170