




EASTERN UNIVERSITY, SRI LANKA
SECOND EXAMINATION IN SCIENCE 1998/99 RE-REPEAT
EXCH 202 MOLECULAR SPECTROSCOPY, AROMATICITY AND REACTION MECHANISM

Time: 02 Hours

Answer **FOUR** questions only

1) Answer **all** parts (a), (b) and (c).

a) Calculate the λ_{\max} values of the UV absorption band of the following compounds



b) i) Give the equation which relates the absorbance of a solution to its concentration and identify all the terms in it.

ii) The UV spectrum of a solution containing 20.5 mg of $\text{CH}_3\text{OCH}=\text{CHC}\equiv\text{CH}$ in 100 ml ethanol when measured in a 2 cm cell had a band at 235 nm with an absorbance 0.70. Calculate the molar absorptivity, ϵ , of $\text{CH}_3\text{OCH}=\text{CHC}\equiv\text{CH}$ at 235 nm.


c) The mass spectrum of $\text{CH}_3\text{COCH}(\text{CH}_3)\text{CH}_2\text{CH}_3$ showed fragmentations at m/e 100, 85, 72, 57 and 43. Give structures of these fragmented ions and indicate the possible pathways for their formation

Contd....

2) Answer **all** parts (a), (b) and (c).

- a) i) The Fieser-Kuhn rule for polyene is given by
 $114 + 5M + n(48.0 - 1.7n) - 16.5R_{\text{endo}} - 10R_{\text{exo}}$
Identify all the terms in it.

ii. Calculate the λ_{max} value of β -carotene.



- b) Explain why acetonitrile (CH₃CN) has resonance at δ 1.97 while methyl chloride (CH₃Cl) has resonance at δ 3.05, even though the electro negativity of cyano group is larger than that of the chlorine atom.
- c) Give the increasing order of C=O stretching frequencies of the following compounds. Give reason(s) for your answer.





3) Answer **all** parts (a), (b) and (c).

- a) An organic compound A (C₈H₈O₂) showed weak absorption at about 3000cm⁻¹, 2850 cm⁻¹ and 2750 cm⁻¹ and strong absorption at 1680 cm⁻¹, 1260 cm⁻¹, 1030cm⁻¹ and 840 cm⁻¹. ¹H NMR spectrum of the compound A had signals at δ 10.0 (s, 1H), 7.5 (dd, 4H), 3.9 (s, 3H).

Interpret the given data and deduce the structure of the compound A.

- b) Sketch the proton NMR spectrum including multiplet patterns expected for the following compounds with TMS as standard. Predict the approximate chemical shifts in your spectrum.





Contd....

- c) Explain why the position of the OH resonance of phenol varies with concentration in solution. But the hydroxy proton of *ortho*-hydroxyacetophenone does not show any great shift upon dilution.



4) Answer **all** parts (a), (b) and (c).

Write the mechanism for each of the following reactions. Indicate all the steps clearly.




5. Answer **all** parts (a), (b) and (c).

- a) By means of equations show how the following conversions may be effected. Give essential experimental conditions.





Contd....

b) Give the structures of the compounds A, B, C, D and identify P, Q, and R.



c) Draw the structures of the products you would expect for each of the following reactions.



6. Answer **all** parts (a), (b) and (c).

a) State Huckel's rule.

- b)
- Use the polygon and circle method to outline the π molecular orbitals of cyclopentadiene and explain, on the basis, why cyclopentadiene is not aromatic.
 - What electron distribution would you expect for the cyclopentadienyl anion?
 - Would you expect it to be aromatic? Explain your answer.
 - Would you expect the cyclopentadienyl anion to be aromatic on the basis of Huckel's rule?

c) Why basicity of aliphatic amines is greater than that of pyridine.

XXXXXXXXXXXXXXXXXXXXXXXXXXXX